
Thermo meter

	Aim :- (34221)
	Him !- (01444)
7	while R To record the body temperature using thermometer
t	Requirement: (3) de 24 ch (1)
1	The amometes
_	a a disaft
).	Refrence: - Right Shanma Shailesh di Gupta G.D Practical of Human anatomy Physiology by Ninali Prakushan (73-75)
	Tractical of Human anatomy Physiology
	by 1412001 tanker shan (13-75)
_	de state de santa de
_	theory: Rteliti) Human are Homoitherms of Nasım bloodid Chemical " the body temperature is maintained between 1705 to 990 f
	Nasım bloodid Chemical
	") the body temperature is maintained between
	* 330t to 330 t
	C
	Heat is lost from body by various mays of Conduction, Convection radication and evaporation C skin, lungs, of excretion
	Conduction, Convection radication and evaporation
	C Skin lungs . It exception
	iv Thalamus Plays an important role in Conservation Cobsorb and loss of heat finam bady, there are Shivering and antrose centers are there in brain that
_	and loss of heat tinom body, there are Shivering
	and antroise centers are there in brown that
-	Sweeting, vasodilation etc. to Inchease heat loss



- The rise in body temporature (pynesia and lever)
 Occurs due to desargement of heat regulating
 mechanisms.
- u) Torins: Poison or harmful Substance Infections, dehydration, tissue etc. also cause use in body
- Temperature in the form of various activities
 like increased metabolism, blood pressure
 pulse, Cardiac output and rate of Respiration
- * A Clinical thermometer is used to measure the body temperature.
- It Consist main bulb and the Capillary tube.

 Because of this device small fortions

 of mercury rises above the twist and

 Separate it self from the main bulk in

 the water bulb.

X

Clinical theirmometer Contains a flattened back with white cooling that helps in seading the index of temperature in certigorade as well as in fatherheit.

OBSERVATION TABLE

S-R	TEMPER	ATURE FROM	BUCCAL	CAVITY	•Ł
1.	99	Temperature	mensure	in ot	1
2.	q 8. V				
3.	97				

SECULA



Procedure:

I Bring the mercury down to about 95° F by quick motion downwords.

- 2) Place it in Such a Position that lissue of Such position one available at compits on axilla, mouth, vogina and rectum. the mouth under the tongue in adults and in compits of Children.
- 3) While taking the temperature under the compit.

 If Should be day Clean and unexpected to the

 Sworounding. No fold of Clothing Should Interface

 the Contact of themometer with Skin.
- 4) While taking the temperature in mouth.

 It should be noted that the patient has not taken

 any hot on Add drink prior to the recording.

 The temperature deported from mouth is always a half degree higher than axillary temperature.

Result: - The body Temperature is found 98. 5%



Aim &- To Record the blood Pressure of a Subject.


John 32)

Reference :- Sharma Shailash & Gupta G.D.
"Practical of Human anatomy Physiology

Requisiement :- STETHOSCOPE, and a mercury SPHYGMOMANOMETER.

Theory: Blood Pressure (B-P) is Defined as the lateral pressure excrted on this walls. of the vessels by the Contained blood. This is Due to the muscularity and elasticity of the walls of blood vessels. The blood Pressure also depends upon the force exerted by the heart When it pumps the blood. The maximum Pressure during systole is defined as systolic blood Pressure Whereas the minimum pressure during diastole is defined as the diastolic blood Pressure. The Difference between the systolic and clicistolic blood Pressure is described as Pulse Ponessure. The average systolic blood Pressure in healthy adult is 100-140 mm Hg.

Waringlight Simulated Mercury Column (LED on LED Display) LCD Standard Cuff Deflation Knob Rubber bulb SPHYGMOMANOMETER



The average diastolic blood Peressone is 80-100 mm Hg.

The average Pulse Peressone is about 30-50 mm Hg.

There are two methods Which are used to
measure blood Peressone using Sphygmomonometer,

Which are ? - (A) Pal Patory method (Feeling Pulse)

(B) Pruscultatory method (Heaving Pulse)

x Procedure:-

- D) Ask the Subject to Sit Down on lie down at ease On the Couch in the Supine Assition, Remove the garments round the armpit, Place the common on a table and adjust Position Such that it is level of the heart.
- 2) The the Cull of Sphygmomanometer and the asim.

 It should be neither to tight to Couse any discomfort


 to the Subject, nor too loose to allow its movement

 yound the com.
- 3) Feel the ordery and mark its Course in the Cabitalfossa.

 also Feel and mark the radial Pulse at the Place
 Well.

3.NO.	SYST	LIC BLOOD P.	
1 : 2 : 3	117	Bb (mmHa)	
-			

JANIFED SECULAR SOCIETA



- 4) Place the manometer by the Side of the Subject. between his own and the body. Tight the Screw of the subber Pump and press the pump to Inflate the bay.

 It inflate to raise the mercury to 200 mm level on 20-25 mmHg Higher after the Disappearance of Pulse.
- 5) Keep the eyes Fixed out merany level and
 The Fingers at the pulse (Which is disappeared)
 Slowly neleasal pressure by unscrewing the Valve
 of the rubber Pump. The reading of the
 level of the mercury when the pulse Proppears
 gives the Systolic Pressure. This is Pripatory
 method this method does not give any idea
 obout diastolic pressure.

Result: - the blood Pressure is found 117 mm tg.

John John

Cate Page

EXPN.3

Aim: To Study-the gross anatomy and Physiology of Digestive Sistem


Reference:

Practical of Human anatomy Physiology by Nivali Prakashan (78.80)


Theory: the Digestive System is a System of body Which bright bound food into Forms that Can be absorbed and used by body Cells. It also absorbs water vitamin and minerals, and aliminates wasks from the body. It bright downs the larger molecules from the body cells by a process Known enough to enter body cells by a process Known as Digestion. The Organs with one briefed in the breakdown of food one Collectively Called the Dogestive System. The digestive System is a tubular System Which extends From the mouth to the anus.

· Mouth and Oral Cavity:

(a) Lips: they are faitly Swammed by Skin and Consists of Striated muscles and mucous glands. The main function of lips is to Penavide a gateury to the objective System.



HUMAN DIGESTIVE SYSTEM



- (b) Teeth: there are total 32 teeth in a normal adult.

 They are further Classified as Incisor (2) Comine(1)

 Premoter (2) and molars (3) On the each Gide of Jaw.

 and Same number of Teeth are there in lower

 Jaw. mayor functions of teeth are Cutting

 Pripping and tearing. Crushing and Gairding.
- (c) Salvary glands: Salivary glands are fresent in an mouth which Secretes Saliva to moisten, lubricate and oligest the food, we engulf. They are of three types: parotid. Sublingual, and Submandibular glands
- (d) Tongue:- Tongue is an accessory digestive system
 Organ Composed of Skeletel Muscle Covered
 With mucous memberane. It helps to taste the food.
 Swallow Food and to Speak. tongue and its associated
 muscles form floor of mouth.
- (e) Pharynx: It is the junction between the Osal and nasal Cavity and inferiorly, it is attached to knynx and Desophagus Pravides lits attachement with digestive System. The major functions of Pharynx areto transport air to the ruspisatory System and food to the digestive System.


(Stomach: Stomach is a I Shaped enlargement of GI Tract Which lies directly inferior to diaphragm.
It Connects desophagus to duodenum (first fart of small Intestine). Stomach Serves as mixing chamber and holding reservoir for food. When food is ingested Stomach pushes asmall quantity of food intoductioner Periodically. As the Size of Stomach can very .
It can Stone large amount of food. It is divided into fown parts: Cordia fundas, Body and Pylowness
The pylowness Communicates with the duodenum of the
Small Intertine via Smooth muscle sphinater called
The fyloric Pyloric sphinater In Stomach, Semi-Sold
bolus its Converted into liquid; digestion of Stanch
Continues, digestion of triglycerides and protein
Storts, and absorption of Several Substances takes place.

(f) Small Intestine: Small Intestine Starts From pyloric sphincter of Stomach, Cails through central and infesior part of abdominal Cavity and at large Intestine. It has major role in digestion and absorption of Nutrients.

Anotomy: It has three major farts:

intestine. It Steats from Pyloric sphincter, extends up to 25 cm and merges into jegunum.

It Secretes proteolytic enzymes (trypsingen,



chymotrypsin, procarboxy, Peftidase, nuclease, Collagenore and clastinase to digest Proteins. It also secretes amylase which digest Carbohydrates and lipase for digestion of lipids.

div Jejunum: It is middle part and extends up to ileum.

It Secretes intestinal juice which is Consists of

Various Proteolytic enzymes like emepsin, arginase


Nuclease etc., Carbotydrates splitting enzymes

like amylase, maltase, lactase etc., and lipid

digesting enzymes like lipase.

(111) Theum: It is last food of Small intestine and ends at fleaceast junction of large intestine. It involves in absorption of digested food material.

Large Intestine: The large intestine is the terminal
Partion of the GI hard. The Overall Functions
of the large intestine are the Completion of absorption.
The production of Certain vitamins, the formation of
feces and the expulsion of frees from the body.
Large intestine is about 1.5 m long and extends from
ileum to amus. The joining of small and large intestine
Occurs at ileacecal sphinder which Controls makement
of material from Small intestine to large intestine.
Large intestine Consists of four major region occum



- (a) Cecum is a small pouch like Organ, which is present next to ile occal sphincler. Attacked to Cecum is a coiled and trusted tube called appendix as Vermisposim appendix.
 - (b) Colon is a long tube which is present next to Cecum.

 The Open end of Cecum attacks with Colon.

 Colon is divided into fown Partions i.e. ascending.

 Glon turns verse colon, descending Colon and Signoid.

 Colon.
- (C) Rectum is approximately last 20 cm of GI tract.
 Tearninal 2-3 cm of rectum is Called anal Canal.
 Opening of anal Canal to exterior is Called anus Which
 is guarded by internal sphincter of Smooth muscles
 and external sphincter of skeletal muscles.
 - The Accessory Digestive Organs: these are the Organs
 Which assists in digestion of food these include
 the teeth tongue, Salivary glands, liver, gall blades
 and pancheas.
- (a) Pancoreas (pan all. Creas Flesh) is a retoroperitored (behind peritoneum) gland, which lies posterior to greater Curvature of Stomach. It is 12-15cm long and 2-3 cm thick. Anatomically, it is divided into three parts;

1. Head: It is expanded position and lies news to conve

2. Body: It is central furt and is left and superior to head

3. Tail: It is last topering fortion of panagers.

These panageratic islets secrete Four types of hormones:

(3) Glucagon: It inconecises blood sugar level.

(3) Insulin: It decorences blood sugar level.

(3) Somato Statin: It mountains Columnagon and Insulin level in body.

(1) Pancacatic Paypeptide: - H Controls Somatostation scretion

(b) Liver and gall bladder:

Liver is the Second largest Organ in body, beded inferior to diaphragm. Gall Hadden is pear shaped sac located inferiorly and Posteriorly to liver. Liver is divided into two lobes: Right lobes (larger) and left tobe (smaller). the inferviour is Called body and the upper tapes postion is Called neck. Liver Produces the bile and the gall bladder stores the bile Secureted by liver and Send it to the small intestine where it aids in the digestive Process.

Result: the human digestive system is studied

C Page

EXPERIMENT NO. 4

Aim: To Study the anatomy and Physiology of


Reference: - Shanma Shrilesh clr. Gupta G.D Practical of Human anatomy Physiology by Ninali Perakashan

Epithedial Tissue: (Epithelial - Guering.) lay on . Epithelial tissue Gonsists of cells arranged in Continuous Sheets in Single on multiple layers. It is living Covering and glandwar tissue of body.

Epithelial tissue Play different roles in body like Protection, Filtration Secretion, absorption and excretion. There are the following three types of epithelial tissue.

I. Simple Epithelialm: It Cordains Single layer of Cells and is of four type:

(a) Simple Epithelium Squamous epithelium: the tissue Consists of a Single typer of for Cells that resembles a tied floor When Viewed from the apical Surface. Simple Squamous epithelium is present at sites. Where the process of filtration are diffusion occurs.





(b) Simple Cubaidal epithelium:

It Consists of Single layer of Cube Shapeol Cells
having sound and Centrally located nucleus.

This epithelium is found in Organs Such as the
thyroid gland and kidney. It performs the functions
of Secaretion and absorption.

(D)

(d).

Simple Columna, epitholium: - the cells of Simple Columns, with Columns, with out nuclei man the base. These play a role in absorption and Scretion. Simple Columns epitholium Cells can be ciliated (containing hair On Swiface.)

Pseudo - Stratified Columnar epithelium:

Pseudo - Stratified Columnar epithelium appears

to have Several layers because the nuclei of the

Cells are at various depths. Even though

all the Cells are attached to the basement

membrane in a Single layer; Some cell do not

extend to the "upper Surface. In Bendo - Statified


Ciliated Columnar epithelium, the cells Secete

mucous or boar Cilia Pseudo - Straitified

non Ciliated Columnar epithelium Conteins

Cells without cilio and lacks mucous Cells.

Cilia Cytoplam Nucleus abstration cell [Diagram of Epitholish They are found in on your skin blood vessels, wantery tract Org



Stratified Epithelium:

It Contains multiple layer of Cells. It is more durable and Can better protect underlying tissues and is of four types:

Co Stratified Squamous epitholium:

Opper layers of this epitholium:

deep core Cubaidal and deeper may be Columnor

This type of epitholium is found at sites

Which receive good deal of abrasion, wear and

tear. Its major functions are Protection,

frevention of water loss and Foreign invasion.

This type of epitholium is generally present in

Skin, Wet living of mouth Desoptagus and tonque.



Stratified Cubaidal epithalium:
This is a fairly rane type of epithelium in which
Cells in the upper layerane Cubaidal, Stratified
Cubaidal-epithelium Serves a role in protection
Secretion and absorption.

Stratified Columnas epithelium:
His also uncommon type of epithelium. the basal layers Consists of Shankned liregularly shaped Cells; Only the upper layor has cells are Columnas in Shape. This type of epithelium functions in Penotection and Secretion.
Peneserd in ducts of glands and Conjuctiva of Eye.

Transitional epithelium:
Transitional epithelium is present Only in the wimay
System. In Its relaxed or Up- Stretched State,
transitional epithelium looks like Stratified
Cuboidal epithelium. As the fissue is stretched
its Cells become flatter. giving the appearance
of Stratified Squarrous epithelium. It allows
the various bladder to Stretch to had a variable
amount of fluid without rupturing.



Glandulas epithelium:

The Function of glandular epithelium is secretion. Glandular Cells often lie in clusters deep to the Covering and lining epithelium. A Gland Consists of a Single Cell our a group of Cells that Secrete Substances into ducts (tubes). Onto a sunface, On into the blood. All glands of the body are Classified as either enlacrine on exacrine.

· Enclocaine glands: (endo-insides crine-secretion)
These are the glands whose secretions (hormones) enter the interstition fluid and then diffuse directly into the blood stream without flowing through a duct. The Pituitury, thyroid and adverse glands are examples of endocrine glands.

· Exocrune glands: - (Exo-Outside; Crine-Secretion) these age the glands whose Secretions (hommones) enter the interstitial stuid secrete their products into ducts that empty ento the Skin Surface are the lumen of abolton organ. the Socretions of exocrine glands includes mucaus, sweet, Oil, ea wax, Saliva and digestive engymes Examples of exocrine glands Include Sudariferous (swed) glands Which Produce swed to help lower body temperature and Salivary. Blands, which Secrete Saliva. Result in the human epithelial Cells are Studied.



[EXPERIMENT. 05]

Aim: To Study the anatomy and physiology of muruka Tissues.

Practice of Human anatomy Physiology by
Ninali Prokashan

Theory: - Muscular lissue Consists of elongated Cells
Called muscle fibres on myocytes that Can
Use ATP to generate force. Muscular tissue
produces body movements, maintains postore and
generates heat. It also provides protection.
Muscular lissue is Classified into three types:
Skeletal, Caroliac, and smooth.

(a) Skeletal muscle tissue: It is named for its
location as it is usually cutached to the bones
of the Skeletan. It Condains Strictions, althorating
light and clark bands within the fibres that one
visible under a light microscope. Skeletal muscle
is Considered Voluntary because it (an be made
to Contact or relax by Conscious Control. A muscle
fibre is roughly cylindrical in shape and has many
nuclei located at the peripheny.



SKELETAL MUSCLE

SMOOTH MUSCLE


CARDIAC MUSCLE

MITED

Diagram of Muscular Tissue



(b) Smooth muche tissue: It is located in the Halls of hollow internal Structures Such as blood vesseles, airways to the lunger the Stomach. Intestines, gall bladder and Orinary bladder. Its Contraction helps in Constriction of blood vessels, physically break objust and I movement of ford along the gastrontestinal tract. movement of fluids through the body and eliminate wastes. Smooth muck fibres are usually Involuntary and they are non- Stricted Clack Strictions). hence named Smooth A Smooth muscle fibre is Small, thickest in the middle and topesing at each end. It Contains a single, Centrally located Cardiac muscle tissue. It forms most of the wall of the heart. It is also Striated. It is Involuntary to Control as its Consciously Controlled. Conding muscle fibres are branched and usually home only one Centrally boated nucleus: an acquisional Cell has two nuclei. They are attached at end to end by interculated discs. Interculated discs hold the fibres together during their vigorous Contraction and Brovide a route for quick Conclustion of muscle action fortentials throughout the heart. Kesufty - the human muscular fissues are Studied



[EXPERIMENT-07]

Aim: To Study the anatomy and Physiology of Nerwus Tissues.


Reference:- Sharma Shailesh Dr. Gupto G.D Practical of Human anatomy Physiology by Mindi Profeshon Page - 53

theory: Nerous tissue Detects changes in Conditions Inside and Ordside the body and sespond by generative action potential (Nerve Impuls) that activate muscless Contraction and glandular Seconetian. Nervous Jissue is Consist of · two types of Cell. Neurons and Heuroglia.

1. Netrons: Newmons are the Cells which are Sensitive to various Stimuli. These Convert Various Stimuli into electrical Signals (alled action Potentials (nerve impulses) and Conduct these action potentials to Other neurons, muscle tissue on glands. Neurons Consist of these

(a) Cell Body: it Contains the nucleus and Other Organelles.

(b) Dendiocites (dends - tree): These are to paing, branched and Short pracesses and are the receiving fortion of a neuron.



(c) Axon: Single, thin cylindrical and long process.

It is the Output portion of a newson which

Conducts nerve impulses toward another newson

or to some Other tissues.

2. Neuroglia: - These Cells do not generate or Conduct nerve Impulses but provide protection and Support to reunans. Neuroglia Cells ere of Six types: astrocytes, Oligo dendrocytes, microglia, ependymal Cells, Schwann Cells and Satellite Cells.

Result: - The human nervous tissue one studied

Exp. No-7 of one's own blood

Réference:— Sterre needle, spêrit, stapes Stopwarch.

Theory: - Bleeding time dépends on sur effectiveness of voisoconstriction and platelet plug jormation. It is deflued as the time from ouset of Breeding till the storage stoppage of bleeding length of time needed for bleeding to cease is noted after dup skin puncture. It determine the function of platelets & integrity of the apillaries. Praeduue July 1118 1181111 million of 11000

1) Clean the finger tip with spirit or alcohol and make a deep puncture with the help of sterich midle.

3) Start Stopwatch Punchlately and Lound the time.

3) Place a deep of Blood on the filter papers after every 30 seconds.

4) Note the time of stoppage of blood.

5) This gives the bledby time.

Result! The bleeding three of my own blood was found to be 3 infinite and 20 seconds.

ing sit is the stand set well the top to the formers on

at zon to produce of the contract

antible de la component de la component de constante de constante de la consta

from to to bush of

Experiencest No - 9
compaire
acué To study of nuccroscope. N: Visture, Practère book of
Reference: Thakare N. Vishuru, Perartère book of Human Anatomy and physiology, Nirali
Human Anatomy and profite 1-4.
Reference: Thakare N. Visheru, Practice book. Human Anatomy and physiology, Nirali Perakashan, Second edition, Page No-1-4.
Reguliements:- nuivoscope
Microscope is an optical l'entrument volviche
uses visible light and a system of linses to
uses morbie again of very small objects.
uses visible itight and a system of very small objects. This unicroscope also called itight unicroscope.
This addrescope
1) compound nuitroscope
2) es simple microscope
3) fleutron microscope
compound ricroscope: It is an awayment of
objeutive and eyepieur use to
majuity an object to the point where it lan
be den with the human eye.
Part of a compound Microscope:
A STATE OF THE STA
1) The support system: The supposed system
constist of various parts which
au disurbed below!
DELTA ® Notebook

a) Base of microscope	
b) Pollans	Page No.
) O Handle (Aum)	
d) Body Tubi	- 15. 1011 H / 12 / 1 (後
e) Stage	7161 30
	wix tuber al
2) The illuminating system: The	ily systems based on
· · · · · · · · · · · · · · · · · · ·	Land Control of the Control
which the microscope function	n de la company
a) Bright field un'croscope.	been last will be
6) Douk field un croscope	No. of the second second
c) & fluorescent uneroscope.	AND PROPERTY OF THE PROPERTY O
d) Condenser.	V 2007
	8. di 2014 1
3) Magnification system: As the	implies majuifying
11 F Cuttury	
I the image of the object wire	al constitution (
compuise of the exprese &	objectives.
a) eye preu: The eyeprees i	the body - Generally
5x and lox experces are used	
manifest of the second of the	11/1/28 1-17/19 12/19/19
b) Objection: - Generally puse	objections and
	u rusolving nosepieu
un a compound interoscope.	U
They a has in a hand the house	Thank in Alle of the Chol
Herachiston appendix:	da chimitud
4) Adjusting system - Two as	yustly system ary
The diffe well in wed;	an the reflection
a) coause adjustment : Two	cause adjusment con
The Bridge of the Screw ow	
course adjustment. These scree	us and mounted
at the top of the hardle	by a double side
micrometer mechanism, on	e on each side.
DELTA® Notebook	

b) From adjustment; Two the adjustment screens and Kejuly - Thu Coarrie mus charitans adjustment screws by doubles side micromatica compounds microscope studied successfully mounted early wide. the handle below the Page No.

compound microscope (Plan paper)

Buaw

Experiment 10- 10 record own pulse note. (se same as experient NO-9) (experiment 9 Requirements: - Stop watch: Throng: The alternating expansion and recoil of elastic autentes during each cardiar cycle creaters a pressure war called a pulse. Thus, the generation of aortic pursure wave that expands the autental wall in seconded as pulse. Pulse Can be felt at vansous antenies ulz, supenficial temporal fanal, common canotid, bronchial, fumoral, potential, modial et .

DELTA® Notebook

L'yocedane;
The same of the state of the same of the s
1) Parpate the superficial auteries by pressing them
assert in adentified bouls.
2) madely at the ward
holling hills Medica
3) Record the pulse mate for 1 min which is expressed
as beats /min.
Result: - My own pulse note was found to be
79 Black/Mills
Contraction of the second of t
Experiment No-11 manual papel of the
Aim: To determine the no. of WBCs un own's blood
Sample.
Remene: Same as exp. No - 9 [Page NO -21-23]
A STATE OF THE STA
Requirements: Hemoustometer Containing Neubauren's
counting country crampen, wis chipette,
steuile needle, nicroscope, coversup, was diluting
fund, spirit in the state of
The state of the s
furniple: - The well of acetic and solo with blood
couse hemolysis of RBCs and accentuates
nuclei of white cells The WBC's com then easily
conted. Mormal large of WBCs in 4500-10000/mm3
of blood the of the state of th
DELTA® Notebook

Theory. The notifie blood cells (WBCs) are also called as leukoustus. They are nucleated, artively amochoid and do not contain harmoglobin and originate purely from extra vascular ussue. They contain proteins and various enzymes. Life span of WBC with the standard of the span of WBCs is less than that of RBCs. The types and subtypes of WBCs: 1) Granwoytes ad all a val Bosinophills is all in 112001 341 a b) Basophill c) Neutrophil 2) Agranulo ytu a) lymphocytes b) Monocytis Procedwee: 1) Take chan and dry apparatus use sterilised needle either having 22 08 24 gauxe number.
Purk the finger and allow blood duop to foun a) Take the pepette and suck the blood bubb pipette up to the 0.5 name only 3) hipe pur tip of the pipette and maintain the blood level at the 0.5 mounts by holding the pipette is a hoursontal position. 4) Dip the pi pette unto un diluting fluid below.

the surface of the regular. suck the WBC diluting fund up to 11 marchs. While the bulb is being filled, top me pipette pipette DELTA Notebook

I with a finger to knock the bead below the surface of fluid in the bulb to prevent any bubble formation. 6) Maintain the level of the mixture at 11 manks by duning unoval of pipette from déluting fluid. 7) Hold the pipette houseoutally and close the both ords of the WBC pipette, then gently mix the contents of the bulb. 8) After mixing, put the pipette un the horizontal Jesition to prevent any loss of its content until 39) Discard the first a duops of fluid from the pipette I as ut does not contain the cells. 3 (0) Charge the entited area of the courting chamber taking came to prevent flooding 311) focus pur nuivoscope conefully and place the 3 Menbaurn's chamber on the stage of the microscope. 12) Look at the country chamber from the Bidi and by turning the course coarse knob, lowers
the power observe (10mm x10mm) with it is
almost touching the cover glass. 13) Now while consinuously looking through the ey's pièce, slowly rotate the coarse adjustment towards yourself will you observe a faint hui in the explicie. Adjust light with a merror ais d'aphragm and use coause adjustment back and fro with you get cells in good focus.

	~~~
Date	
Page No	

Requirement of Automorphisms of RBG.

Requirement of Autoscope, himocytometer contactuly

Requirement of Autoscope, himocytometer contactuly

Newbarrent's counting chamber, RBG

pripette, sterils needle (32-24 gauge), caverslip

RBG distriby fluid

Theory: The rad blood cells also called as

engthrougher, have chicalan, biconcaut

slape. They are non-nucleated cells and

Page No. are red in colour. Their lifespan is of 120 days. formation of RBCs un bom manufur is known as enytheropoiesis. The nain function of RBCs is to. of the body and carbon de oxide from cells / Hissues - The Momal Hange of RBCs are 4.5-6 millions mm3 of blood. OVER THE BUILDING OF THE STATE Puocedure o 1) Take furthey purposed RBC deluting fruid un a I watch glass. 2 2) Steurise the skin on the finger with a cotton 3 swah dipped in spirit and putick with steurised meedle. 3) Suck blood unto the pipette up to 0.5 work on 4) Hold the pipette un nouixonal position and by planly tip of pipette against index finger and blowing gently on the nubber tube, adjust the blood column, so that it is exactly on the o.s work and suck the deluting fund immediately up to 101 mark of the pipelte. 5) Rotate the pipette jou 2-3 numertes with its of the contents of the tube. The med bead in the bulb of the pipette should move from the one state to other during the nutring of blood cells.

6) Discound the first two drops of fluid from the bipette, which probably does not contain cells. 7) Placed lep of the pipette on the edged on sented

DELTA Notebook

Page No.

auso and below one two duops of diffully blood

3) Place the danged chamber on the star of microscope
and odjust the microscope for observation water low

powers (tox)

9) Focus the central square of the Meribaner's
chamber and dark the uniform distribution of fels.

(o) There forms the REC square explan high power

Objective (40 x). count in the medium REC square

8). Country of REC should have from small equares

at upper left hand torner

b) A cell that faither any of the trible thus

that border the right star of REC Seiten grand

hat be counted.

(3) count the cell of clark paper of the seiter grand

REC count, has penjourned by mising of blood with on the hours on the land to be conserved on factor to be conserved that

RB(s count = Total no. cells un 5 RBC sections X

10,000 correction factor (200 x56)

= Xe elocoo

Finithions 1 mm 3 of blood

Result The total RBC Count of own blood was

found to be smillion 1 mm 3 of blood.

four to be 5 million mind of bloods which the the sould like Experiment No-13101 with the To determine the blood quoup Alu = Build saulples all parting a share the saulples Same as Requirement. - Glass stides or stide nelde 3-4 caucties, Steulle needle, spirit, autiseum A (Blue), B (yellow), C (Colonules). 00000000 ABO Bliod guoup is based on two glycollpiols
housely autigen A and Andiger B. Peuson neith Antigen
A on their RBCs feel under blood geoup A, those
with B antigen on their RBCs have blood
group B and those who have Both Ap and B
antigen on their RBCs feel under blood Theory :-DELTA® Notebook

an houses neither Ac	8
Thought and Thoughtans with passes	
B antigens full under blood geroup o and in	- 原
Kuow ii au minerisa acres	
blood to only penson.	T.C.
AND DEFEND DESCRIPTION OF THE STATE OF THE S	12/20
Puocedure & De material de la	
The state of the s	
1) Crean and dry the carrity wide.	
2) Mark the affiles of the Slide as H, B, D.	
3) Place of auti A, Anti B, and and B	1年。
ses berefully all the calley.	
4) steurise finger lip with a cotton small dipped	
un sphit and puck the finger using a steal	ised
needle.	
51 Add 1-2 duops of blood us each depuession	and
mixed ut Immediately with the help of a sealed	
glass capellary or by gentle shaking of carrity	
stide on plain houixava strujare	15
6) observe agustration journation in can depress	ion.
Carlotte and the second second	100
Result: - The Blood group of my own's blood Sam	10.
was found to AB.	
The state of the s	NY.
The state of the s	25
The second secon	
The state of the s	A TOP TO STATE OF THE PARTY OF
The state of the s	
The section of the se	
All step and specific to the state of the st	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
And the same of th	
The state of the s	
The state of the s	

Experiment No-14 To study the candiovasculari system (CVS) The ball of the °- 87 7 93 ( Page Sauce as - candivasculan (cus) consist of heart, blood (1 1 A FL 10 and bluod nessels. a connective tissue composed of a liquid Blind is extracellular matrix called blind plasma that contains dissolved substances and contains vanious cells bragments suspended win it 1. Blood transport og e nutrients. Blood flows from heart towards blood bessels or from blood vessely to hour . This function is counted out by uts pumping actions s want in actsulas vas pumpinning, willing plusia appround on Minh is through Anatomy of hearth - 18 , miles what a day were our The total Jud in the state of th (A) location of the heart of The mant we dies blue Veutebrou, Column and lung. cixe and chape of heart, on It is 12 cm, long , 9 cm heide 6 cm thick neith about 250-300 gm weight with proposited and their world with pulse but 2 C) Layers of heart: + pulandium * epilardium. Myocaudium. * Endocaudium. DELTA® Notebook

I w u chamber,
D) Heart chambers: To is composed of 4 chambers,
two atura superior
Ventricus infectionly.
20 - 1 - 1 - 1 - P. I m
* Right ventricle.
1
* Left verdusch.
Hel verdeich .
Systemic and pulmorary circulation:
Systemic Circulation: This involves flow of
oxygenalated blood from left
ventucle unto aorta turough the St valves, then
into antenies, outtenibles and finally and
systemic capillaures in which exchange of gasts
and metrients take place. Blood unloads by
and takes up Con though capillatures and venules
and the same travel towards right attitum
of heart from velu and the process continues.
Pulmorary circulation = During Venturellar diastole
deoxygenated blood from the right
ventulcle flows unto the pulmonary trunk, which
branches into pulmorary autericis that carry
blood toward the lugs.
The state of the s
Conduction systems
The second of th
Heard beats due to its inherent and ellything c
electrical activity. Autorhythmus februs acc
responsible for such actually, they repeatedly
DELTA® Notebook

9 11 5	
The state of the s	
0 1 5	Page No.
12 3	generale au aillon potential du to volvich the
2 11	heard unisdo.
21-3	
1-1-5	Autourhythung fibers: These are as paremaker and four
J-11-2	the conduction system - a specialised
-1.5	network of condear muscle liber, gives direction for
The state of the s	each eyell of caudige excitation to progress through
1-1	Pull 14 1/1 1 1 / KI I WI I I I I I I I I I KI I I KI I I I
1-3-	a) SA Node
05	b) Dy Woods
- 4 0	c) Burdle of his.
	d) Burdle of Branches
- 3	e) Punkinje fibrus.
- 3	
3	Candiae agele:
3	
- 11 3	Events occurry during each beat of heart is
	called as caudian Lycle. Alormal Rate 72 beats/
N-C-EURI	
	during each condicae cycle.
	* Atrial Systole
	* Ventucular Systoli.
P - 3	* Ventricular Bastoli
	31000 vessels: - Blood vessels and of fine types
3	consisting of antendes, antenides,
- CC	aprillancies, venules and venus.
30	will: The cus are was studied successfully.
	The cost of Successfully

exp-15

Aim

To study the human respiratory system. Page 10 (103-106)

The functioning of the respiratory system along with the circulatory system is essential for the exchange of oxygen and carbon dioxide (gases). Respiration is the chemical process in the biological system where in exchange of these gases take place (cellular respiration is the constant supply of oxygen to cells for energy-conversion process that occur in the mitochondria of the cells). The exchange of gases between blood and the lungs is called external respiration, while the exchange between blood and the cells is called internal respiration.

Anatomically, the respiratory system has two parts viz. upper respiratory tract (nose, nasopharynx, oropharynx, laryngopharynx and larynx); lower respiratory tract (trachea, bronchial tree including, bronchioles, alveolar ducts, alveolar sac and alveoli) (figure 26.1)

Physiologically, respiratory system includes various accessory parts, oral cavity, rib

cage and respiratory muscles (diaphragm).




Figure 26.1: Respiratory tract

# Upper respiratory tract

- (a) Nose: It is divided into external and internal structures. External structure consists Nose: It is divided into external and hyaline cartilage which surrounds the muscle and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and supporting framework of bone and hyaline cartilage which surrounds the muscle and support framework of bone and hyaline cartilage which surrounds the muscle and support framework of bone and hyaline cartilage which surrounds the muscle and support framework of bone and hyaline cartilage which surrounds the surrounds th supporting framework of bone and nyamassing the outer side of each oval cartilaginous expansion forming and supporting the outer side of each oval nose lies on the analysis opening is called as the ala. The nasal passage of internal nose lies on the roof of the mouth. The nasal passage is separated into right and left cavity by a middle portion called as the septum. In side the nasal cavity, the frontal region is known as vestibule The nasal passage consists of pseudo stratified ciliated columnar epithelium cells. The olfactory receptors (for smell sensation) are situated on lining of the superior  $n_{asa}$ conchae and near the septum. The functions of the nose include-warming of inhaled air and entrapment of particulate matter and bacteria in the sticky mucous.
- (b) Pharynx: It is also known as throat. It is funnel shaped, up to 13-14 cm long and lined by mucus membrane. Fibrous and mucus tissues begin at the internal half and extend to the cricoids and inferior cartilage of the larynx. The pharynx lies posterior to the nasal and oral cavities, superior to the larynx and just anterior to the cervical vertebrae and forms the common passage for food and air, and provides a resonating chamber for voice production. It is divided into nasopharynx, oropharynx and laryngopharynx.
- (c) Larynx: The larynx is known as the voice box, located between the root of tongue and upper end of trachea and in front of the lower part of the pharynx. The shape of larynx is triangular with attachment of cartilages to one another by ligaments. The cavity of larynx extends from its triangle shaped inlet at the epiglottis to the circular outlet at the lower border of the cricoids cartilage, where it is continuous with the lumen of the trachea. The pairs of folds of larynx form 3 compartments viz. vestibular, vocal and lower true vocal cords. The slit like gap between the true vocal cords is called as rima glottidis. Glottis is the space between the rima glottidis and true vocal cords.
- (d) Trachea: The trachea or wind pipe is a tubular structure, 11-12 cm long, extends from the larynx in the neck to the bronchi in the thoracic cavity and divided into right and left bronchi. The major layers of the tracheal walls are mucosa, submucosa, hyaline, cartilage and adventitia. The 16-20 incomplete, horizontal rings of 'C' shaped hyaline cartilage provide a semi rigid support and hence prevent the collapse of tracheal wall during inhalation.

104

- (e) The bronchi: The right and left bronchi begin at the level of the 5th thoracic vertebra. The right bronchus is vertical, shorter and wider that the left one. Carnia is the internal ring, from where trachea divides into right and left bronchus. The mucus membrane of the carnia is the susceptible part of the larynx and trachea involved in cough reflex. After entering into the right lung, the right divides into 3 branches while the left bronchus after entering into the left lung, divides into two branches, one goes to each lobe of the lung. Each branch then divides into bronchioles and subdivides into smaller bronchioles. Bronchi are lined with ciliated columnar epithelial cells and the terminally bronchi are lined with simple cuboidal epithelium. The bronchioles are divided into alveolar ducts, and these ducts result in the formation of bag like structures known as alveoli which are responsible for the exchange of gases. The main functions of bronchi are adhering to foreign particulates and bacteria due to their sticky nature and the movement of mucus towards the throat.
- (f) The lung: The lung is a cone shaped organ located in the thoracic cavity, that extends from the diaphragm to a point slightly above the clavicles. Heart and other adjacent organs in the mediatstinum, separates the thoracic cavity into structurally different chambers. Pleural membranes protect the lungs from any damage. The superficial layer called the parietal pleura, lines the wall of the thoracic cavity, deeper layer known as visceral pleura, covers the lungs. Pleural cavity is the space between the parietal and visceral pleurae which contains a lubricating fluid, which aids in reduction of friction between membranes. Due to the space occupied by the heart, the left lung is about 10 % smaller that right lung. The main and important function of lungs is the exchange of gases.

#### Physiology of respiratory system

Respiration is a process by which the lungs inhale the air in and thereby expand the size and contract to exhale air.

### Phases of respiration

When the capacity of the thoracic cavity is increased by contraction of intercostal muscles and diaphragm, pressure in the thoracic cavity decreases as compared to the atmospheric pressure. These results in air being drawn into the lungs to equalise the alveolar and atmospheric pressure in the process called inspiration or inhalation.

When the diaphragm and intercostal muscles relax, the ribs regain their original and this results over position, the diaphragm ascends, and the lungs recoil and this results expiration of the diaphragm ascends. exhalation. There is a brief pause (stoppage of breathing) and another cycle begins.

# Pathological significance

Pharyngitis: It is the inflammation of pharynx.

Laryngitis: It is the inflammation of larynx.

Asthma: It is a disorder characterised by obstruction in airway, inflammation, hypersensitivity which causes difficulties in breathing. These symptoms may be due torelease of chemical substances, histamine, bradykinin and leukotrienes (inflammatory mediators) in response to antigenic stimuli.

A

T

the m

urine stora

with

const

To

U

Chronic obstructive pulmonary diseases (COPD): These diseases are characterised by persistent airway obstruction due to emphysema or bronchitis.

Chronic bronchitis: Chronic bronchitis presents itself as cough with or without expectoration for at least 3-4 months of a year for 2 consecutive years.

Emphysema: It is the enlargement of alveoli with destruction of alveolar septa.

Pneumonia: It is an infectious disease characterised by alveolar inflammation, fever, chills productive or dry cough, malaise (pain throughout body) and chest pain Streptococcus pneumonia is the common causative microorganism responsible for the pneumonia but it can also be caused by viruses, fungi and parasites.

Tuberculosis: It is an infectious and communicable disease caused due to Mycobacterium tuberculosis affecting lungs and pleurae of bronchi, and it is characterised by symptoms mainly fatigue, loss of weight and appetite, cough, chest pain etc.

RedeRatory

### **EXERCISE**

- What is respiration? Differentiate between external and internal respiration.
- Enlists the organs of upper and lower respiratory tracts.
- 3. Give the anatomy of bronchi.

	Date: 12 Phonon 12017.	Mariano de Carros de Carro
	Experiment -15	Nas
Ain	To Study winery system with the help of model and charte.	6
V.	Reference - Waugh Anne and Grant willon hous and wilson, Anatomy and Physiology, 11th edition; churchill living stone Publication. Page no 332-344-	10
-	Anatomy and Physiology, 11th edition; churchell alling seems	
-	Requiement → charte and modeli.	
	Theory -	
Property of the second	The uninary system is the male excretory system	
<u> </u>	and consut of:-	
(d)	and consist of!—  2 Kidney which secrete where.  2 weter which convey wine from kidney — wingry  bladder.	
	The univary bladder, where wine collects and a tempor	la=
9	illy stille.	
	bladder to the excreter externor.	
-1	Hunction 3-1	and the second s
	Detailitation - It helps un memoral of tours from the body.  Homeostand It helps to maintain homeostasin by sector  Teacher's Signature	
<b>D</b>	Homeostario de helps to maintain homeostario by rector	ug

porsal acta Pest canal Adrenal body. - umay duct -urinary bladdel -wall of bladder cut off to show openings of urinary duct Exactory agans

Teacher's Signature .....

	Date . 1 120	
	er demoring selecture amount of water or electrolyte un	The state of the s
-1	Kilney - Et ill bean shaped, 10-12 cm dong and weight	
	auaund 150 gm. Et is clocated b/w the partietal ID specietonium and posterios spect wall of the abdomen	
	he dayen of where that surrounds each kidney are !-	
	The dayen a white that suspring each kidney	The second second
	aue!-	-
0	Enner dayer - Renal caprule.	
(2)	Middle dayer - Adipore caprule.	
<b>3</b>	Outer Mayer - Renal Jaseia.	
7	priorital section of kidney decieals ar outer reddich	
	area - corper and unner reeddlik brown area - meder	
~ ~	frontal section of kidney vieweals ar outer reddeit area — cortex and unred reddeith brown area—medu. Medulla consist of conical shaped striations, the renal housement.	
	ckeramids.	
		1
-	Mephron - It ill the smallest functional and structural unit of Kläney.  Et consults of aux poutsons -	The second secon
	unit & Klåney.	A STREET, SQUARE, SQUA
-	Et consiste of aux fourtions -	-
$\mathcal{O}$	Renal Capule:	
(2)	Renal Capsule: Genal Publile:	
	Renal capsule - It will composed of and components-	
2	glomenulus and the glomenilus capsule.	
2))	Renal tubule - It conside of - purinal concertated orbito	•
	glomenulus and the glomenulus capsule.  Genal tubule - It consid of -proximal convulated tutule,  Jook of Henle and diffal convulated tubule.	1

Pyranua of medullary sutitione kenal artery heral vein - contex PeluD Lis of Kidney

Tunction of nephron-2) Filtration. The upper part of the nephron in - senal cortex and the down faut (400) of Henle in the Iseral medulla. - Functione of Ridney-Formation and secretion of wine.

Rusduction and secretion of engtheropoletin.

Buduction and secretion of menin. A. - Genal Tubule -The dutal communated trubule cleading unto a collecting duct which write forming fakulary duct that empty with the mion cary. The cells of proximal consulated tubule are subordal aplithelial cells and have a forominent buidge border of micho-villi- it sincueaux othe surface area you weaksoftion and secretion.

In accending don't and full faut a accending don't flence, the cells are double aguarded

Page No. 30

Date: / /20

epithelial colls. The thirt accending dumb of Henle is composed of simple cuboldal to columnas epithelial cells. itue. - Junction - Enwlued in reabsorption and secretion. - Tirata Glomerular apparators ne macula derra celle along ultr juxta glomerular celle your juxta glomerulus apparatu - Formation of wine - It stakes three steps for wine formation -O Letteration - Boumanns Capsule, glomerulus. Describe dealsosption- Publical part 3 Secretion - collecting duct, distal chibile. B. - Jenal capsule - one close end with the wenal tukule form the cup shaped Boromann's capsule which almost completely encloses a network of anternal capillaries.

Called domesulus. Called glomerulus. Boiomani capiule in a double layered structure consulting of unceral layer and familial layer. → Function → Pruolued ûn the gelteration of wire. Uneter 35-30 cm dong, 3mm un diameter.

consult of Gruee dayers?

Outen dayer country of aerolas connective true

contanous with fibrious capsule of the kidney.

Teacher's granture.

middle layer in a varcular dayer composed of unner itual dayer of muscle fiker

Enner clayer al a mucour membrane with transitional all epithelial and assolar connective triure with consider able amount of collager and elastic fixure and elymphatic itime.

The weter in continous with Junnel Shaped weral pelvis.

Function - They are the buber that carry wine from the kidney to the winary bladder t

Huethra? The weethra is a canal extending from the to mech of the bladder to the exterior at the external weethral evifice. It is donger in the female.

The male unothers un auxited with the unrary and

The yemale unethua un approximately 4cm long and 6mm un diameter. It une downwards and fourant behind the symphysic fueld and from at the external unetheal auffice just in the front of vagina.

The external wrothval arrive in quarded by the external wrothval sphinoter which is under itue udurtary canal central. Hrnary Bladder - Ristansible muscular augar schrated un feluic causty. It can expand to hold about 700-800 mi of wine. The bladder wall its composed of three Outer dayer of doore connecture deine containing blood, blood versels, neure and symphatic verseli. Middle dayer - consult of three dayers of smooth mucle -fibre - inner dongitudinal, middle circular and outer dongitudinal mude dayer his us alled detruser Prince Mayer - of transitional opethelium. The three orifices 4 in the bladder form a bladder wall strangle. The Upper two are the opening of mother, down in the your of origin of weether. Smooth muscles of bladder wall were down unto weether where the weether originate the circular fibre form an internal sphendure (gate like structure) that controls the passage of They from the bladder Below the unternal sphinder il the external sphinder which il compared of skeletal muscle. The untrinal sphlnoter all under autonomic neuau contro system (ANG). Function of Kidney >

Page No. 83 Regulation of blood wolume and work composition

(electrolyte balance).

Maintainence of blood osmolarity.

Regulation of blood puessure.

Regulation of blood ph.

Regulation of blood glucore devel.

Excuetion of waste and fourign substance. iture mat k Result - The excuetory system was studied with the help of models and character.

# EXPERIMENT NO. 39

20

# STUDY OF THE HUMAN SKIN

exp: 17.

> Aim

To study the human skin

Some (

Theory

Skin is part of the integumentary system which helps in protection and temperature regulation. It is also known as cutaneous membrane, an outer covering of body with surface area of about 1.5-2 m² and consists of glands, hairs and various cells. Skin has two important layers i.e. epidermis and dermis (Figure 39.1).




Figure 39.1: The Human Skin

**Epidermis :** It is the superficial layer of the skin, made up of stratified keratinized squamous epithelium which varies in thickness in different parts of the body. Blood vessels and nerve endings are absent in the epidermis, although for supply of oxygen and nutrients, deeper layer are bathed in the interstitial fluid of the dermis.

The keratinocytes, melanocytes, Langerhans and markel cells are the important cells found in the epidermis. Keratinocytes produce a protein called as keratin which helps in protection from heat, chemicals and microorganism. Melanocytes are cells which produce a pigment known as melanin which gives a characteristic colour to the skin and protect skin from UV light. Lagerhans cells originate from the bone marrow, migrate towards the epidermis and prevent the entry of microorganisms into the skin. Merkel cells formed deep in the epidermis detect touch sensation. Epidermis is made up of stratum basale, stratum lucidum, stratum spinosum, stratum corneum, stratum granulosum.

Stratum basale is made up of cuboidal columnar ketatinocytes, which form a tough protein keratin which helps in protection from injury to deeper layer. Stratum spinousm superficial to stratum basale gives strength and flexibility to the skin. Stratum granulosum, a middle portion of epidermis is made up of 3-5 flattened keratinocytes, the keratinocytes acts water repellent sealant, retard loss and entry of water and foreign materials. During apoptosis (programmed cell death), break down of nuclei take place, and thus keratinocytes of stratum granulosum do not carry out metabolic reactions and they die. Hence, stratum granulosum marks the trasition between the deeper, metabolically active strata and the dead cells of the superficial strata. Stratum lucidum found in fingertips, palms of hands and sole of feet provides toughness. Stratum corneum is the outer layer of epidermis offers protection to microorganisms and injury.

Dermis: It is a tough, elastic layer made up of dense irregular connective tissue collagen fibre. The different cells found in the dermis are fibroblasts, macrophages adipocytes along with blood vessels, nerves, glands and hair follicles. Dermis is divided into papillary and reticular region. The papillary region is made up of collagen and elastic fibres, and perceives the sensation of pain, warmth, cold, tickling and itching, whereas reticular region consists of fibroblasts, macrophages, and adipocytes. The network like arrangement of collagen fibers helps the skin resist stretching. Collagen and elastic fibers provides the skin with strength, flexibility and ability to stretch.

Sweat glands: These are distributed throughout the skin, predominantly found in the palms of hands, soles of feet, axillary and groin regions. Tiny pours of sweat gland secrete clear, watery fluid; due to decomposition of bacteria it has unpleasant smell.

Hair: These are present on skin surface except on the palms and soles of feet. Hair is divided into shaft (above the skin) and root (portion that penetrates into the dermis and use

to the subcutaneous layer). Hair follicles are the part surrounding the root and made up of epithelial root sheath. The onion shaped structure at the base of each hair follicle is known as bulb, which contains a nipple shaped serration called papilla, made up of areolar connective tissue and blood vessels responsible for nourishment for hair follicles.

Sebaceous glands: These are found in the skin of the scalp, face, axillae and groin region. They secrete sebum into hair follicles and thus keep the hair soft with shiny appearance and prevent infection against bacteria and fungi. Sebum also prevents drying and cracking of the skin. During puberty, there is excessive secretion of sebaceous glands rendering the skin prone to the effects of excessive moisture.

#### Functions of skin

- 1. Provides protection against microorganisms and UV light.
- 2. Regulates body temperature by maintaining homeostasis
- 3. Synthesis of vitamin D through exposure to UV light;

4. Prevent dehydration due to moisturizing action.

away studited

#### **EXERCISE**

- L. Enlist the layers of skin.
- 2. Discuss in short the structure of skin.

exp:-18

> Aim

To study the human pectoral girdle and bones of the upper limb

Theory

### Pectoral Girdle:

The pectoral girdle or shoulder attaches the bones of the upper limbs to the skeleton. There are two pectoral girdles consisting of clavicle and scapula.

Clavicle: It is an 'S' shaped bone, also known as collar bone. It is positioned horizontally across the frontal part of the thorax, superior to the first rib (figure 29.1). It has convex and concave curves, and the junction between these form the delicate region of the clavicle most likely to fracture. It articulates with manubrium of the sternum and forms the sternoclavicular joint. Clavicle articulates with scapula at acrominoclavicular joint and with humerus at the shoulder joint. The clavicle transmits force from the upper limb to axial skeleton and also provides attachments for the muscles.





Figure 29.1 : Clavicle bone or Collar bone

Scapula (shoulder blade): It is flat and triangular shaped, located on the back surface of the chest wall and is superficial to the ribs (figure 29.2). From the lateral side, it has shallow surface known as glenoid cavity which is involved in the formation of shoulder joint with the head of the humerus. Spine is a rough ridge on the posterior part of the body which extends beyond the lateral border of the scapula and overhangs the glenoid cavity. The acromion process with clavicle forms the joint called as acrominoclavicular joint, a slightly movable synovial joint responsible for the movement of shoulder girdle. The coracoid process, is a projection from upper boarder portion of the scapula that provides attachment to muscle and gives movements to shoulder joint.



#### Figure 29.2: Scapula

#### Bones of the upper limb

Bones of upper limb consist of humerus, radius and ulna, wrist bones, bones of hand, finger bones etc.

Humerus: It is the longest bone of the upper limb. The head of the humerus fits into the glenoid cavity of scapula, forming the shoulder joint also called as socket joint. The greater and lesser tubercles are the two rough projections of humerus. There is a deep groove present between these two tubercles known as bicipital groove (figure 29.3). The distal end of the humerus presents two surfaces that articulate with the radius and ulna to form the elbow joint.

for

ca

(fig

ne

th

W

pa

th

d

fi




Figure 29.3: Humerus

Radius and ulna: The radius is the smaller bone and situated on the lateral side of the forearm. Its proximal end is narrow, having disc shaped head which articulates with capitulum of the humerus and the radial notch of the ulna, where as the distal end is wider (figure 29.4). Adjacent to the head of the radius is the neck, and roughened area inferior to neck is the radial tuberosity, which is a point of attachment of tendons of biceps muscle. At the distal part of the radius, there is wider shaft forming a styloid process on the lateral side which gives attachment to brachioradialis muscle and radial collateral ligament to wrist part of the hand.

The ulna is located on the medial part of the forearm and is larger than the radius. The olecranon, the proximal end, forms the prominence of the elbow. An anterior projection to the olecranon is the coronoid process which attaches with trochlea of the humerus. The distal end of the ulna consists of head which is separated from the wrist by disc of fibrocartilage. Posterior to ulna's distal end there is styloid process, which gives attachment for the ulnar collateral ligament to the wrist. Both the radius and ulna form the elbow joint with the humerus.




Figure 29.4: Radius and ulna

Wrist (carpal) bones: The eight carpal bones, closely fitted and arranged into two row i.e. proximal (scaphoid, lunate, triquetrum, pisiform) and distal rows (trapezium, trapezoic capitates, hamate). The proximal bones of carpal are associated with the writ joint wherea distal bones form the joints with metacarpal bones.




Figure 29.5 : Carpal, metacarpal and phalanges

Metacarpal bones: The fives bones of the palm of the hand are called as metacarpal bones. The proximal ends articulate with carpal bones and distal ends with phalanges.

phalanges: There are 14 phalanges, 3 in each finger, and 2 in thumb. Phalanges articulate to metacarpal bone and with each other by hinge joint (figure 29.5).

### Thoracic cage

Thoracic cage consists of bones of a sternum, 12 pairs of ribs and 12 thoracic vertebrae




Figure 29.6: Thoracic cage

Sternum: It is also known as breastbone. It is flat, narrow, 'T' shaped, lies in the middle of the anterior thoracic wall and has 3 parts; manubrium, gladiolus and xiphoid process (figure 29.6). It connects to the rib bones through cartilage, forming the anterior section of the rib cage with them, and thus helps to protect the lungs, heart and from physical trauma.

Result: - The human bettoral gledle and bonce of the upper White was

# **EXPERIMENT NO. 30**



# STUDY OF BONES OF THE VERTEBRAL COLUMN

To study the bones of the vertebral column

Theory

The vertebral column is known as backbone or spinal column and with the thoracic cage it forms the skeleton of the trunk of the body. The major functions of the vertebral column are to facilitate the forward, backward and sideways movements. It serves as a point of attachment to the thoracic cage and muscle, protects the spinal cord and provides support to the head.

There are a total 26 vertebrae which include 7 cervical vertebrae at the neck region; 12 thoracic vertebrae posterior to thoracic cavity; 5 lumbar, 1 sacrum and 1 coccyx. The sacrum and coccyx are the immovable bones while others are movable.

#### > Cervical vertebrae

A total of 7 cervical vertebrae smaller than thoracic vertebrae make up 3 foramina i.e. 2 transverse and 1 vertebral foramina having vein and nerve fibers passing through.

Atlas: It is the first and topmost cervical vertebra, and along with the axis, it forms the joint connecting the skull and spine. Atlas is a large ring like structure without body, consisting of an anterior and a posterior arch and two lateral masses.




Figure 30.1: Atlas Vertebra

The superior articular facets of atlas articulate with occipital conclyles of the occipital bone (figure 30.1) to form the atlanto-occipital joint which gives the 'YES' saying movement (up and down movements) of skull. The inferior articular facets articulate will, axis.

Axis: It is the second cervical vertebra, and has a strong odontoid process which rises perpendicularly from the upper surface of the body (figure 30.2). It forms the pivot joint by articulating with atlas which helps during the rotating of head side to side.

The other 4 cervical vertebrae consist of bifid spinous processes and 7th is non-bifid spinous and serve as essential anatomical landmark.




Figure 30.2 : Axis

#### Thoracic vertebrae

Thoracic vertebrae are 12 in numbers which are larger and stronger than other vertebrae. It articulates with ribs to form facets (figure 30.3),




Figure 30.3: Thoracic vertebra

It consists of heart shaped body, downward pointed spinous processes, two facets present, on either side, two costal semi-facets, one above, near the root of the pedicle, the other below in front of the inferior vertebral notch. A small facet at the tips of the transverse process articulates with tubercles of the ribs.

## The lumber vertebrae

They are 5 in number and is the largest vertebrae of the column. The body is kidney shaped and does not articulate with ribs (figure 30.4). The spinous processes of this vertebra are flat and stout and have backward projection and provide attachment for the back muscles.




Figure 30.4: Lumbar vertebra

### The sacrum

The sacrum is a triangular bone formed by the collection of five sacral vertebrae which articulates to the 5th lumbar vertebrae and forms the lumbo-sacral angle and below with the coccyx (figure 30.5). The body is large and much wider from side to side than from backwards. Its anterior projecting edge is named the sacral promontory. Transverse process and the costal element are fused to each other to form the upper surface of the lateral mass of the sacrum on both sides. Its base projects forward and upward forming the prominent sacrovertebral angle when articulated with the last lumbar vertebra. Its central part is projected backward, so as to give increased capacity to the pelvic cavity. In between the

body and lateral masses contain anterior sacral foramina which offer gap for passage of nerves. The inferior surface of the sacrum articulates with coccyx.





Figure 30.5: The sacrum

## The coccyx

It is triangular, single, fused with 4 rudimentary vertebrae. Superiorly, it articulates with sacrum and contains a series of transverse processes on the lateral surface (figure 30.6).



# Figure 30.6: The coccyx Bones of Verytebrai 1. Enlist the bones of the vertebral column

- 2. Give the locations of the thoracic vertebrae.
- 3. What are the functions of the cervical vertebrae?
- 4. Give the total number of war a

# STUDY OF VARIOUS JOINTS

Exp- 20


, Aim

To study the various human joints

Theory

Joint is a point of attachment or contact between two bones or between bones and cartilages, or teeth. Joint is also called as articulation.

joints are classified into



### 1. Fibrous Joint

Sutures

Syndesmoses

Interosseous membrane

#### 2. Cartilaginous joint

Synchondroses

Symphyses

### 3. Synovial joint

Ball and Socket joint

Hinge Joint

Pivot joint

Condyloid joint

Saddle joint

Plane Joint

### 1. Fibrous joints

Fibrous joint holds the other articulating bones closely together by dense irregular connective tissue and is without synovial cavity; thus these joints do not move or show little movements.

**Sutures**: It consists of thin irregular connective tissue that join the bones of the skull logether by its interlocking edges (figure 32.1). The interlocking edges offer strength, and reduces chances of fracture and also acts as shock absorbent.

**Syndesmoses:** These are fibrous joints, articulate two bones and are arranged as ligaments which permit limited movements of bones (figure 32.2).




Figure 32.1: Suture ligaments

Interosseous membrane: It consists of dense irregular connective tissue which binds the adjacent bone and allows slight movements e.g. interosseous membrane between radius and ulna; tibia and fibula (figure 32.3).




Figure 32.2 : Syndesmoses of tibia and fibula




Figure 32.3: Interosseous membrane between tibia and fibula

### 2. Cartilaginous joints

Cartilaginous joints permit slow movements or no movements due to absence of synovial cavity. The bones of these joints are tightly joined by hyaline cartilages.

chondroses: It is a cartilaginous joint in which the epipyseal plate of each end of a one connects the epiphysis of growing bones by either hyaline cartilage or tilage (figure 32.4).




Figure 32.4: Synchondrosis

symphyses: The ends of articulating bones are covered with hyaline cartilage and nect the bones. e.g. the pubic symphyses found in symphysis between the frontal aces of hip bone (figure 32.5).




Figure 32.5: Symphysis

#### . Synovial joints

It consists of synovial cavity which offers movements of joints and thus these joints are freely movable. The cartilages of these joints are smooth and slippery for free movements.

Ball and socket joint: It consist of depression or cavity on one side of a bone and ball like surface on the another bone, this arrangements permits them to fit closely to each other and allows free movements. The shoulder and hip joints are the examples of ball and socket joint (figure 32.6).




Figure 32.6: Ball and socket joint between acetabulum and femur

**Hinge joints :** These joints permit unidirectional movements e.g. elbow joint, interphalangeal joint (figure 32.7).




Figure 32.7: Hinge joint between humerus and ulna (at elbow joint)

Pivot joints: The round surface of the bone attach to another bone partly by ligaments and allow the movement around its longitudinal axis. e.g. atlanto-axial joint (figure 32.8)




Figure 32 .8: Pivot joint between head and radius and radial notch of ulna

Condyloid joints or ellipsoidal joints: It is the convex oval shaped bone fused into depression of another bone and permits two sided movements e.g. radiocarpal and metacarpophalangeal joints (figure 32.9).




Figure 32.9: Condyloid joint of radius and lunate and scaphoid bones of carpus (wrist joint)

Saddle joint or seller joint: In this, one bone is saddle shaped and the other bone is used into the saddle. It allows the two sides movements same as that of ellipsoidal joints. g. carpometacarpal joints (figure 32.10).




Figure 32.10: Saddle joint between trupezium of capus and metacarpal of thumb

### Significance

Rheumatism: It is the disorder of joints and connective tissues characterised by pain, inflammation and difficulty to initiate movements.

Osteoarthritis: It is a group of mechanical abnormalities involving degradation of loints, including articular cartilage and subchondral bone characterised by pain, lenderness, stiffness, and locking.

Rheumatoid arthritis: It is an autoimmune disease that results in a chronic, systemic inflammatory disorder that may affect many tissues and organs. It is characterised by inflammation of joints, swelling and loss of function.

Ankylosing spondylitis is a chronic inflammatory disease of the axial skeleton with variable involvement of peripheral joints and nonarticular structures. It is characterised by pain and stiffness in the hips and lower back that progress upward along the backbone.

RUMIH


Joeley

wey

Studled

### EXERCISE

- 1. Define a joint. Give the classification of joints with examples
- 2. What are fibrous joints? Enlist joints of the fibrous trans-



The CNS: The main function of CNS is to maintain integration of sensory input and motor output, thus it enables a person to identify sensory information and formulate exponses to changes that alter haemostatic balance. Apart from this function, the nervous stem is also involved in perceptions, behaviours, memories, and all voluntary movements. The nerves (neurons) are a bundle of thousands of axons plus associated the modern to the control of the contr

he CNS consists of brain and spinal cord

The Brain: The brain lies within the cranial cavity, and constitutes 1/5th of the body leight and consists of cerebrum, brain stem (mid brain, pons), medulla oblongata, and lefebellum (figure 33.1).




Figure 33.1 : The Human Brain

Cerebrum: It is the seat of intelligence and is the largest part of the brain and occupinanterior and middle part of the cranial cavity. It is derived by a deep cleft into right and le cerebral hemispheres. Inside the brain, the two hemispheres are connected by corp callosum.

Cerebral cortex is the superficial part of the cerebrum. Hemisphere of cerebrum  $h_{as}$  lobes viz; frontal, parietal, temporal and occipital lobes.

**Function of cerebrum :** Mental activities are involved in memory, intelligence, sense responsibility, thinking, reasoning, moral sense and learning.

- Sensory perception (pain, temperature, tough, taste, smell)
- Initiation and control of voluntary muscle contraction

  Functional areas of cerebrum
  - 1. Motor area
  - 2. Sensory area
  - 3. Brain Stem
- **(b) Mid brain :** It is situated around the cerebral aqueduct between the cerebrabove and pons varolii below. It connects the cerebrum with lower parts of the brain a spinal cord.

- (c) Pons varolii: It is situated in front of the cerebellum, below the mid brain and above the medulla oblongata. It consists of nerve fibers forming a bridge between the two hemispheres of cerebellum and of fibers passing between the higher level of brain and the spinal cord.
- (d) Medulla oblongata: Extends from the pons varolii above and is continuous with the spinal cord below. It is pyramid shaped and it lies within the cranium above the foramen magnum. Its anterior and posterior surfaces are marked by central fissures. Cells of the medulla oblongata consist of relay stations for sensory nerves passing from spinal cord to cerebrum.

Vital centers present in medulla oblongata are given as below.

- 1. Cardiac centre controls functioning of the heart.
- 2. Respiratory centre controls rate and depth of respiration.
- 3. Vasomotor centre controls the diameter of blood vessels.
- 4. Reflex centre for vomiting, coughing, sneezing, swallowing.

**Decussation of pyramids:** Motor nerves descending from the motor area in the cerebrum to the spinal cord in the pyramidal or cortico spinal tract, crossing from left side to the right and vice versa in the medulla oblongata.

2. Cerebellum: The second largest part of the brain, situated posterior to pons varolii and immediately below the posterior portion of the cerebrum lying posterior to cranial fossa. It has two hemispheres, separated by narrow median strip called as gray matter which is found on the surface, while white matter is present inside the cerebellum.

Cerebellum performs functions like coordination of voluntary movement, posture, balance and coordinates activities associated with maintenance of balance and equilibrium of the body.

### Spinal cord

It is elongated and almost cylindrical part extending from the brain just below the medulla. It is suspended in the vertebral canal and is surrounded by three meninges and cerebrospinal fluid

Meninges are: 1) dura matter, 2) arachnoids and 3) pia matter.

The spinal cord extends from the first cervical vertebra to the lower border of the first lumber vertebrae, is approximately 42-45 cm long and is about the thickness of the little finger (figure 33.2).




Figure 33.2 : Spinal cord

The spinal cord is the nervous tissue link between the brain and the rest of the body, except the cranial nerves. Motor nerves originating from the brain descend through the spinal cord and supply to various organs and tissues at the appropriate level of the cord. Sensory nerves from different organs and tissue enter and pass through the spinal cord to the brain. This sensory information is conveyed by the dorsal nerve route and spinal cord. The cell bodies of these unipolar, sensory neurons make up a small region of gray matter in the dorsal root ganglia; while fibers of the ventral nerve root carry motor information out of the spinal cord. Cell bodies of these multipolar, motor neurons are in grey motor area that composes the inner core of the spinal cord.

Diencephelon – extends from the brain stem to the cerebrum and surrounds the third ventricle.

It comprises of the thalamus, hypothalamus and epithalamus.

The thalamus has important structures, which relay sensory impulses from the sensory areas to the spinal cord and brain stem. The thalamus receives, relays, and transmits sensory and motor information to and from the cerebral cortex.

Inferiorly to the thalamus is the hypothalamus having a number of nuclei; those involved in various behavioral and physiological functions.

lyron Ichiai

# Important functions of the hypothalamus are given below

Regulation of ANS functions like contraction of smooth muscle and cardiac muscle; production of hormones, regulation of emotions and behaviour, food and water intake, equilibrium, control of body temperature and circulating circadian symptoms and consciousness.

Epithalmus is superior to thalamus and contains pineal gland which secretes melatonin involved in induction of sleepiness.

The arrangement of grey matter in the spinal cord resembles the shape of the letter H, having two posterior, two anterior and two lateral columns. The area of grey matter lying transversely is the transverse commissure and it is pierced by the central canal, an extension from the fourth ventricle, containing cerebrospinal fluid.

The white matter of the spinal cord is arranged in three columns or tracts; anterior, posterior and lateral. These tracts are formed by sensory nerve fibers ascending to the brain, motor nerve fibers descending from the brain and fibers of connector neurons.

There are two main sources of sensation transmitted to the brain via the spinal cord.

- 1. The skin: Sensory receptors (nerve endings) in the skin, called cutaneous receptors, are stimulated by pain, heat, cold and touch, including pressure. Nerve impulses generated are conducted by three neurons to the sensory area in the opposite hemisphere of the cerebrum where the sensation and its location are perceived.
- 2. The tendons, muscles and joints: Sensory receptors are nerve endings in these structures, called proprioceptors, which are activated in response to stretch phenomenon. Together with impulses from the eyes and the ears they are associated with the maintenance of balance and posture and with perception of the position of the body in space.

Motor neuron stimulation results in contraction of skeletal (striated, voluntary) muscle, contraction of smooth (involuntary) muscle, cardiac muscle and the secretion by glands controlled by nerves of the autonomic nervous system.

### Spinal Cord Physiology

It provides conduction pathways to and from the brain and serves as the integrator or reflex center for all spinal reflexes. The white matter tracts in the spinal cord are highways for nerve impulse propagation. The grey matter of the spinal cord receives and integrates incoming and outgoing information.

### Pathological Significance

Stroke: It is a neurological disorder due to blockade of blood vessels which supply blood to the brain tissue. Symptoms include paralysis of one part of the body (hemiplegia). loss of sensation, slurred speech, death may also occur.

Predisposing factors include hypertension, atheroma, cigarette smoking, diabetes mellitus, blood clot, high cholesterol in blood vessels, stress etc.

Dementia: It is a permanent or progressive loss of intellectual abilities i.e. impairment of memory, judgment, obstruct twinkling etc.

Meningitis: Inflammation of meninges due to infection caused by bacteria or virus. Symptoms include fever, headache, vomiting, confusion, lethargy and drowsiness.

Epilepsy: It is the disorder of CNS characterised by periodical and unpredictable occurrence of seizures. Seizures occur because of rhythmic, synchronic discharge of neurons due to hyper activity leading to behaviour impairment with or without affecting the motor activity. Depending upon the factors and part of the brain involved, it is of two types, generalised (throughout brain) and partial (local).

Various antiepileptic drugs used in the treatment include- Benzodiazepines, barbiturates, valproic acid, phenytoin etc. alone in combination for certain period.

Depression: Depression is a feeling of intense sadness or unhappiness. It develops due to illness, hormonal imbalance, environmental factors, genetics etc.

Management of depression can be done by psychotherapy and drug treatment was studied (imipramine, fluoxetine etc). system Rosull -

EXERCISE

Give the main functions of the CNS.

- Enlist the various structural parts of the brain.
- Give the functions of hypothalamus. 3.
- Give the physiology of spinal cord. 4.
- Define the following term
  - Depression



# EXPERIMENT NO. 34

# STUDY OF THE LYMPHATIC SYSTEM

exp 22

(25)

Aim

To study the lymphatic system

Theory

The lymphatic system is a part of the circulatory system consisting of a network of, lymphatic vessels for transportation of lymph. Lymphatic tissue is a reticular connective tissue that contains a large number of B and T cells (lymphocytes). Lymph enters via vessels of increasing size into a large number of lymph nodes before returning to blood circulation. The lymphatic system helps in circulating body fluids and in defense against diseases causing agents

### Lymph and lymph capillaries

Lymph is a clear, pale yellow fluid (in intestine it is creamy white), similar in composition to that of plasma and interstitial fluid and varies in plasma protein content. It transports the lymphocytes, plasma proteins, bacteria and cell debris of damaged tissues which is filtered and destroyed by lymph nodes.

Lymph capillaries are composed of a single layer of endothelial cells that originate as blind end in the interstitial space. They are more permeable to proteins and cell debris. The minute capillaries join together to form larger lymph vessels which cover the arteries and veins. Lymph vessels consist of numerous cup shaped valves that ensure the unidirectional flow of lymph towards the thorax. The contraction of adjacent muscles and regular pulsation of larger arteries induce milking action which facilitates the onward movements of lymph. The thoracic and right lymphatic ducts are formed by the association of lymph vessels together and they empty the lymph into the subclavian veins.

Lymph node: Lymph nodes are bean shaped, located along lymphatic vessels in a group. It is 1-25 mm long, covered over by a capsule of dense connective tissues that extends into the node, forming the trabeculae, which divide the node into compartments. Lymph node contains lymphocytes and macrophages. Nearly about 4-5 afferent lymph vessels enter the lymph node and only one efferent vessel carries lymph from the lymph

node. Hilum is the concave surface of the lymph node, where in the artery enters and values. Internal to the capsule is a supporting network of reticular fibers and fibrobiac which along with capsule and trabeculae constitute the stroma of the lymph node, the parenchyma of lymph node is divided into a superficial cortex and deep medulla, the superficial cortex consists of outer and inner cortex. The egg shaped B cells known as lymphatic nodules are found in outer cortex, while the inner cortex contains T cells and dendritic cells that enter the lymph node from the other tissues. The medulla contains a cells, antibody producing plasma cells and macrophages.

## > Functions of the lymph node

- 1. Lymph node acts as a filter and traps foreign particles;
- 2. Phagocytosis i.e. engulfment of particulate matter, bacteria, cells from malignary tumor;
- 3. Multiplication and proliferation of T and B lymphocytes

#### > Spleen

Spleen is oval shaped, and is the largest lymphatic organ ,12 cm in length, 2.5 cm in thickness and contains reticular and lymphatic tissue (figure 34.1). It lies in the left hypochondriac region between the stomach and diaphragm. Impressions are the notches found in the adjacent organs of lymph which includes gastric impression (stomach), renal impression (left kidney), and the colic impression (left colic flexure of large intestine). Similar to lymph node, spleen also has hilum, through which splenic artery, vein and efferent lymphatic vessels enter. The capsule, trabeculae, reticular fibres and fibroblasts form the stoma; the functional part i.e. parenchyma of the spleen consists of red and white pulp. Red pulp consists of a dense network of fine reticular fibres, continuous with those of the splenic trabeculae, to which are applied flat, branching cells. The meshes of the reticulum are filled with blood. The removal of ruptured, worn out, or defective blood cells and platelets by macrophages, storage of platelets and blood cell production during foetal life are the important functions carried out in the red pulp of the spleen. The white pulp consists of lymphocytes and macrophages. The blood circulates into spleen via splenic artery as it enters the central arteries of the white pulp.

B and T cells perform immune functions and macrophages are involved in phagocytosis in white pulp of spleen.




Figure 34.1: The spleen

### > Thymus gland

in

S,

6

It is located anterior to the sternum and posterior to the heart having weight of approximately 30-40 g in adults. It consists of 2 lobes attached by areolar tissue. The lobules of thymus consist of framework of epithelial cells and T lymphocytes. Thymus gland is involved in development and production of mature T lymphocytes. The maturation of T lymphocytes is initiated by hormone called as 'thymosin'.

## Pathological significance

Lymphoma - the cancer of the lymph nodes

Hypersplenism - abnormal splenic activity because of enlargement of spleen associated the excessive destruction of blood cells.

Red Results Lymphon

**EXERCISE** 

system was studies

- I. What is lymph?
- . Give the role of lymph capillaries.

# STUDY OF THE MALL " experior 23

> Aim

To study the male reproductive system.

Reproduction is a process in which an animal or plant produces an individual like itself.

The male reproductive system consist of external genital organs viz, scrotum, penis and urethra and internal reproductive organs such as testes, vas deferens, seminal vesicles and prostate gland bulbourethral gland etc.

Scrotum: It is a pouch like structure hanging from the base of the penis (figure 35.1). The scrotal septum divides vertically into two sacs, each one containing testes. The position of scrotum and contraction of its muscle fibers regulate the temperature of testes.

Testes: Testes are two in number, suspended in each sac of scrotum by spermatic cord. Each testis is about 5 cm in length, 2.5 cm in diameter with 10-20 gm in weight and covered by tough fibrous capsule. Each testis contains spermatogenic cells and sertoli cells in somniferous tubules, supporting the process of spermatogenesis. Somniferous tubules produce sperms, which during intercourse, enter the vagina and into fallopian tube of female reproductive system where it fertilises the ovum (released from ovaries) in the fallopian tubes. Testes are involved in spermatogenesis and secretion of male sex hormones i.e. testosterone from interstitial or Leydig cells.

Penis: It is a male genital organ for reproduction consisting of urethra for excretion of urine as well as semen ejaculation. It is cylindrical in shape and consists of glans penis, and the body. The body of the penis is composed of fibrous tissue known as tunical albugenia. The dorsolateral muscles are called the corpus cavernosa. Upon dilation of blood vessels of corpus cavernosa, erection of penis takes place causing maximum blood to enter into blood sinus (figure 35.1).

Epididymis: It is a 4 cm long, comma shaped organ that lies along the posterior border of each testis. It is divided into head, body or tail. The tail continues as vas deferens. It is the site for sperm maturation, where the sperms acquire motility. Epididymis is also the storage site for sperms and helps to propel them by contraction of vas deferens.




Figure 35.1: Male reproductive system

Vas deferens: As the tail of epididymis continues, it leads to vas deferens, which is about 45 cm long; dilated part of vas deferens is called as ampulla.

Vas deferens stores and expels the sperm during ejaculation by contraction of muscular wall.

The urethra: It forms the terminal part of the duct of the reproductive and urinary system, and is the common pathway for semen and urine

The seminal vesicle: It is the accessory sex gland of the male. It consists of two pouch like structure, 4-5 cm in length. Each seminal vesicle secretes a viscous alkaline fluid which contains fructose for ATP production by spermatozoa. Each seminal vesicle communicates to form ejaculatory duct. Fluid secreted by seminal vesicle normally constitutes about 60% of the total volume of semen.

Prostate gland: Prostate gland is located in the pelvic cavity anteriorly to the rectum. The prostate gland secretes milky alkaline fluid, which aid in lubrication during intercourse. It composed of citric acid, proteolytic enzyme and constitutes about 25% of lotal volume of semen.

Bulbourethral gland: This gland situated inferior to prostate on the either side of membranous urethra. It secretes alkaline fluid into urethra in sexual stimulation. It also secretes mucus that helps in lubricating the penis during intercourse.

### Pathological significance

Cryptorchidism: It is a condition in which testes remain at undescended position in the inguinal ring, sometimes in abdomen. Crytorchidism may occur due to short spermatic cord, narrow inguinal canal, adhesion to peritoneal or inadequate testosterone secretion. Cryptorchidism may result in infertility, inguinal hernia and malignancy.

Elephantiasis: It is the thickening of scrotum skin which looks look elephant's hide, leading to enlargement of scrotum.

Prostatitis: It is the inflammation of prostate gland occurs commonly due to bacterial infection mainly through urethral route. Result I

Kepuedutiu

### **EXERCISE**

- Enlist the various male reproductive organs with their functions.
- Cive the anatomy and -

# EXPERIMENT NO. 36

# STUDY OF THE FEMALE REPRODUCTIVE SYSTEM

, Aim

To study the female reproductive system.

, Theor

Female reproductive system consists of ovaries, fallopian tubes, vagina, vulva (external genital organs) and breasts (figure 36.1).




Figure 36.1: Female reproductive tract

Vagina: It is a tube lined structure, 10-12 cm long that begins from the cervical region of the uterus. It functions as a passage for the foetus at the time of delivery, menstrual blood during menses and sperms during sexual intercourse. A recess called fornix is attached to the cervix on which the intrauterine devices (contraceptive device) especially diaphragm are placed. Vaginal smooth muscles adequately stretch during child birth and intercourse. The superficial layer of the vagina, called the adventia is composed of connective tissue. At the vaginal orifice, there is a thin membrane called hymen, forming border of the orifice (os) that get ruptured during intercourse.

Vulva: It includes the following structures-

Mons pubis: It is anterior to the urethral and vaginal openings and is composed of adipose tissue, covered with skin and pubic hair.

Labia majora: These are two folds extending from mons pubis, to the perineum. It is made up of adipose tissue, sebaceous glands, and sweat glands with pubic hair.

Labia minora: These are two folds on both side of the vagina composed of fatty tissue, sebaceous glands and sweat glands.

Clitoris: It is located anterior to the labia minora. It is a cylindrical mass of erectile tissue. The stimulation of clitoris produces sexual excitement in female.

**Vestibule :** It is the space between the labia minora (inner labia or vaginal lips, flaps of skin on either side of the vaginal opening and is surrounded by labia majora) and is occupied by the vaginal orifice, the hymen and the urethral orifice. This tissue helps to narrow the vaginal orifice during sexual intercourse which puts pressure on the penis. Within the urethra and vaginal orifice are the paraurethral mucus secretion and bartholins gland respectively. These glands secrete mucus during sexual excitement and aids in lubrication during intercourse.

Breasts (mammary glands): This gland is responsible for synthesis and expression of milk under the influence of hormone viz; prolactin and oxytocin. Each breast has nipple, behind which are present the lactiferous ducts and sebaceous glands. Pigmented area around the nipple is the areola .The breast is supported by supensory ligaments, which pass through the skin lying deeply inside the breast. Each breast has 20-25 lobes separated by adipose tissue .The lobules of breast contain alveoli and is surrounded by myoepithelial cells. Contraction of these cells under the influence of oxytocin (a hormone released from the posterior lobe of pituitary gland) results in the expression of milk. Lactation is the process of synthesis and ejection of milk from breasts.

Ovaries: These are two in number which produce sex hormones (estrogen and progesterone) and the ova. Each ovary lies in the shallow cavity on the lateral wall of the pelvis attached to uterus by broad ligament called as mesovarium. Each ovary measures about 2-4 cm in length, 2 cm in breath and is 1.20 to 1.40 cm thick, consisting of fibrous tissue, blood and lymphatic vessels, nerves etc (figure 36.2).

Fallopian tubes: These are two in number, lying one on each lateral side of fundus of the uterus. Each tube measures about 10 cm long. These tubes are responsible for transport of ovum towards the uterus. The funnel shaped region of fallopian tube is known as in infundibulum. The infundibulum has finger like projections. The fallopian tube is made up of three layers of viz. serosa, muscularis and mucosa which help in the movement of the

ovum. Also due to peristaltic action of the muscularis, ovum is propelled along




Figure 36.2: Ovary

Uterus: Uterus is present between the bladder and rectum. It is the site for implantation of fertilised ovum, development of the embryo and foetus and also a passage for menstrual bleeding. It is a pear shaped organ having 7.5 to 8 cm long, 5 cm wide and 2.5 cm thick having fundus, body and cervix. Uterine cavity is the space between the uterine walls. Uterus also has three layers, outer layer is perimetrium, myometrium (middle having smooth muscle) and endometrium. Myometrium produces coordinated contraction under the influence of oxytocin during labor, helps expulsion of foetus from the uterus. Endometrium has two important layers, stratum functionalis and baselis. Stratum functionalis sloughs off (rupture) at the onset of menstruation while stratum basilis gives rise to formation of new stratum functionalis after menstrual bleeding.

# Pathological significance

Dysmenorhea: Pain during menstruation occurs due to tumour, formation of ovarian cysts, pelvic inflammatory disorder, intrauterine devices, oral contraceptive usage etc.

Pelvic inflammatory disease: Bacterial infection of pelvic organs (uterus, uterine tube, ovaries) characterised by pelvic soreness, lower back pain, abdominal pain etc

Vulvovaginal candidiasis: It is characterised by itching, thick yellowish vaginal discharge, pain, and commonly occurs due to fungal infection with Candida albicans. Antifungal drugs like Fluconazole are commonly used for treatment of candidiasis infection due to Candida albicans.

Polycystic ovarian syndrome (PCOS) : This syndrome is characterised by oligomenorrhoea, anovulation, infertility, hirsutism and obesity in young female with enlarged and cystic ovaries. This may be due to excess androgen secretion and less follicle stimulating hormones.

Dysfunctional uterine bleeding: Excessive bleeding during or in between menstrual periods without a causative uterine lesion. It occurs commonly in association with anvolutary cycle i.e. menarche (beginning of the cycle) and menopause (cession of menstrual cycle). Anovulation occurs due to hyper secretion of estrogen alone without progesterone.

Premenstrual syndrome: It is cystic disorder characterised by edema, weight gain. breast swelling, pain, constipation, anxiety, irritability, headache, etc. A complete and det with carbohydrates, proteins, fibre and analgesic drugs (ibuprofen or diclofenac sodium) can reduce the severity of this disorder.

Endometriosis: It is a condition in which cells from the lining of the endometrial layer grow outside the uterine cavity and is characterised by symptoms such as bleeding, severe pain during menstruation and infertility. The main sites for abnormal endometrial development are ovaries, uterine ligaments, rectovaginal septum, pelvic peritoneum.

Result !-

### **EXERCISE**

- 1. Enlist the various organs of the female reproductive tract
- 2. Describe the anatomy of the ovary
- 3. Define the following terms
  - (a) Endometriosis,
  - (b) Dysmenorrhea,
  - (a) Dalum

# **EXPERIMENT NO. 37**

STUDY OF THE HUMAN EYE

exp: 25

Aim

To study the human eye.

Theory

The eye is a special organ for the sense of sight. It is located in the orbital cavity; it is spherical in shape and 2.5 cm in diameter. The eye ball measures about 2.5cm diameter. The orbit of the eye is formed by different bones which protect the eye ball. Anatomically, the wall of the eye ball is divided into 3 layers. Outer layer is a fibrous coat; consisting of sclera and cornea, the middle layer is a vascular coat consisting of choroid, ciliary body and iris. The inner layer known as retina consists of nervous tissue (Figure 37.1).




Figure 37.1: The Human Eye

Sclera: The white region of eye is made up of collagen fibers and fibroblast. The
fibrous tissue of sclera maintains the shape of the eye and also provides site for
attachment to the extrinsic muscles of the eye.

- 2. Cornea: It is 0.5-1.0 cm thick, made up of epithelium, it functions in transmission light toward retina as well as provides effective barrier against bacterial invasion.
- 3. Choroid: It is situated posterior to the sclera and is very vascular, deep chocolate brown in colour. Light enters the eye through activating the nerve ending in the retina and finally absorbed by choroid.
- 4. Ciliary body: Presents anteriorly to the choroid, consists of the non striated muscle fibers, epithelial cells and contains ciliary process. Secretion from epithelial cells forms aqueous humor. The size of pupil changes due to contraction and relaxation of ciliary muscles under postsynaptic stimulation and occulomotor nerve.
- 5. The iris: It is anterior to ciliary body, between the cornea and lens, consists of circular and radial smooth muscles. At the centre of the iris is the pupil; the size of the pupil changes during contraction of circular muscle and radial muscle. Circular muscle has parasympathetic innervations and upon activation it contracts (meiosis) and the size of pupil reduces; this is observed in bright light. In bright light, the sympathetic nerve innervates the radial muscle and its activation leads to relaxation of radial muscle (mydriasis).
- **6. Lens :** It is a circular- biconvex transparent body. The suspensory ligaments of the ciliary body holds the lens enclosed within the capsule. It is elastic in nature and lies posterior to the pupil.
- 7. **Retina**: It lines in the inner surface of the choroid and is composed of nerve cells fibers. They acts as rods and cones and are activated depending upon the intensity of light passing into the eye. The vitreous humour in the eyeball cavity, posterior to the lens, helps in maintaining intraocular pressure to support the retina against choroid.

# Physiology of vision

Photoreceptors and photo pigments – Stimulation/ activation of the photo receptors takes place when the image of the object is focused on retina, thereby transfering light to bipolar cells. The rhodopsin of rod and other pigments of cones are responsible for colour vision.

Optic tracts: It is the pathway posterior to the optic chiasma. The tract consists of nasal fibers of retina. The tract goes back side to meet lateral geniculature into the center for visual information received from the retina of the eye. It collects information

the ascending retinal ganglion cells through the optic tract and from the reticular activating system and thus neurons sends their axons via optic radiation.

## Pathological significance

Cataract: It is the clouding of the lens inside the eye which leads to decrease in vision.

It is the most common cause of blindness and is conventionally treated with surgery.

Glaucoma: It is a group of ocular disorders with multi-factorial etiology, characteristic with increase in intraocular pressure-associated optic neuropathy that results in impairment invision.

Cycloplegia: It is paralysis of the ciliary muscle of the eye, resulting in loss of accommodation.

#### **EXERCISE**

- 1. Give the anatomy of the human eye.
- 2. What is the role of retina?
- 3. Give the physiology of vision.
- 4. Describe the following

# **EXPERIMENT NO. 38**

# STUDY OF THE HUMAN EAR

> Aim

To study the human ear

> Theory

Ear is the organ of hearing. The cranial nerve (8th cranial nerve) carries sensation of hearing towards the cerebral cortex which is a region for the integration and interpretation of information.

Ear is divided into 3 regions

External Ear: Collects sound waves and transmits towards middle ear.

Middle Ear: Middle ear also called as tympanic ear, which conveys sound vibrations to the oval window

Internal Ear: Receives and interprets the sound for hearing and equilibrium.

External ear: Consists of the following parts.

**Pinna OR Auricle:** It is the expanded part, made up of fibroblastic cartilage covered with skin. The expanded outer edge is known as helix and lower most part is the lobule. Ligaments and muscle attach the pinna to the head.

**External auditory canal :** It is an irregular shaped long tube about 2.5cm in length which lies in the temporal bone and leads from auricle to the ear drum. It is divided into cartilaginous and osseous part. Cartilaginous part is about 1/3rd of auditory and contains ceruminous glands which secrete wax called cerumen that helps to trap the entry of unwanted particles into the ear.

Middle ear: It is small and filled with air and is known as tympanic cavity. The middle ear contains a chain of three small bones called auditory ossicles of which one end is connected to the ear drum and other end to fenestra vestibuli. Auditory ossicles also contain malleus, incus and stapes (figure 38).

The handle of the malleus attaches to the internal surface of the ear drum. The hand of the malleus articulates with the head of stapes. The base of the stapes fits into the oval window.

**Internal ear:** Internal ear is bony and in the form of a membranous labyrinth. The space between two labyrinths contain perilymph fluid, whereas, membranous labyrinth contains endolymph fluid.




Figure 38.1 : The ear

The bony labyrinth is divided into 3 parts

Vestibule is an expanded part of the middle of ear contains fenestra vestibule and cochlea.

The cochlea is snail shaped organ having broad base and apex. The three canals are continuous with the vestibule. The membranous labyrinth in the vestibules consists of sac called utricle and saccule which are connected by small ducts.

At one end of each semicircular canal is a swollen enlargement called the ampulla. The vestibular branch of the vestibulocochler nerve (13th cranical nerve) consists of ampullary, utricular and saccular nerves. Cell bodies of the sensory neurons are located in vestibular ganglia. Auditory nerve goes posteriorly through temporal bone and reaches the hearing area of the cerebral cortex.

### Physiology of hearing

The auricle or pinna collects the sound waves and transmits them into the external auditory canal, where it strikes the ear drum. Because of the change in the pressure of air, the ear drum vibrates. The central area of the ear drum connects to malleus, which also vibrates. These vibrations are transferred to incus and in turn to stapes. As the stapes

vibrate, the membrane of oval window is pushed in and out. The oval window vibrates about 20 times more vigorously than the ear drum because of transmission of small vibration oscillates over a larger surface area. The movement of oval window sets up fluid pressure waves in the perilymph of the cochlea. As the oval window bulges ahead, it pushes the perilymph of the scala vestibuli. The scala vestibule transmits pressure waves to the scala tymphani and then to the round window, causing it to bulge outward in the middle ear building pressure waves in the endolymph ( rich in potassium ions) present inside the cochlear duct. Vibration of basilar membrane in the endolymph, allows hair cell movement of spiral organ against tectorial membrane. Bending of stereocilia ( mechano sensing organelles of hair cells )leads to receptor potential which results in the nerve impulse formation through auditory nerve to hearing area of the cerebral cortex where the interpretation of wave take place.

### Pathological significance

Deafness: hearing impairment, or hearing loss is a partial or total inability to hear.

Otitis: Inflammation of ear due to infection.

pes aly 1

#### **EXERCISE**

- 1. Enlist the various parts of ear
- 2. Give the anatomy of the ear.
- 3. Give the physiology of hearing.
- 4 What is deafnoss?

# RECORDING OF ELECTROCARDIOGRAM

Ex10-27

> Aim

To record Electrocardiogram (ECG)
Requirements

Electrocardiograph with leads

> Theory

Electrical current generated in the heart by action potential can be detected/recorded by ECG on the body surface. This will record the signals generated by heart muscle fibers during each heartbeat. The ECG is recorded by the use of chest leads and limbs leads. These leads positioned on the chest, arms and legs, amplify electrical signals and thus produce various tracings that enable the determination of conduction pathway, size of heart, causes of chest pain and areas of heart damage etc. The instrument used to record these changes is called as an electrocardiograph.

#### Procedure

- 1. Ask the subject to lie down and relax.
- 2. Apply the jelly at left and right wrist and left and right leg above the ankle joint to hold the leads.
- 3. Connect the electrodes at the above mentioned positions.
- 4. Switch on the machine and keep the stylus at the centre of the paper.
- 5. Adjust the sensitive to get standard.
- 6. Adjust the lead select or knob to record the ECG.
- 7. Place the chest electors in proper position on the chest after cleaning and application of jelly.
- 8. Record the ECG and observe various intervals as shown in figure 16.

Following tracing can be observed in normal ECG

P Wave - Deflection produced by atrial depolarisation

QRS complex - Represents rapid ventricular depolarisation. Impaired or abnormal conduction of heart results in widening of QRS complexes

Wave- Indicates ventricular depolarisation

ST Interval-Represents the time when the ventricular contractile fibers contract during plateau phase. ST intervals increase during myocardial infarction and reduce in coronary ischemia, hypokalemia and digitalis toxicity.

QT Interval - Represents the electrical depolarisation and repolarisation of the left and right ventricles. Prolongation of QT interval is observed in ventricular tachyarrhythmias and shortened QT can be associated with hypercalcemia.




Figure 16.1: ECG graph

#### Pathological Significance

- 1. The use of ECG is used to analyse the relative size of the heart
- 2. Study of distribution of rhythmical quality and conduction system of heart Diagnosis of myocardium ischemia, pericarditis caused due to Mycobacterium infection and hyperkalemia (high K⁺)

Result

**EXERCISE** 

1 What is FCG?

> Aim

To measure th

Theory

Body temper

safe and effective healthy body thermoregulatio varies, although accompanying controlled cher

Technique

thermostat set i

- Oral c reliable
- 2. Tympo emiss audite
- 3. Axill them

facto

4. Rect

	Alem.	
S.NO.1	Proceedie	
L	To second the body temp using the theumometers	in the second
2 3	To Herord the Blood pressures of a subject of To study the gross anatomy and physiology of	
44	To study the anatomy and physiology of apithelial tissues.	
805	To study the anatomy and physiology of musuland	
6.	To study the androwy and physiology of Newvous	
g. 0	To determine the bleeding time of one's own blood. To determine the blood clothing there of ours own	
9· 00	To study of microscope.  To Record own bulse mate: (Pres No- 65)	
11 12 13.	To distribute the no. of Wes un own's blood so	1.5%
140	To determine the Blood group of one's own Blood so To study the Condiovascular system.	xmple.
15,	To study the wuran webbatown	
	Out Collin King	
19	To study the human performs giridle and	
1		
1	o study the various human joines	
10	study the human central memory system	
	study the lymphath system.	

Frankali

To study the funde reproductive system.

To study the burnan eye.

To study the burnan eye.

To study the burnan eye.

To study the burnan eye.